

# The Central Limit Theorem

Patrick Breheny

March 6

## Kerrich's experiment

- A South African mathematician named John Kerrich was visiting Copenhagen in 1940 when Germany invaded Denmark
- Kerrich spent the next five years in an interment camp
- To pass the time, he carried out a series of experiments in probability theory
- One of them involved flipping a coin 10,000 times


# The law of averages

- We know that a coin lands heads with probability 50%
- Thus, after many tosses, the law of averages says that the number of heads should be about the same as the number of tails ...
- ...or does it?

## Kerrich's results


Number of tosses ( $n$ )	Number of heads	Heads - $0.5 \cdot$ Tosses
10	4	-1
100	44	-6
500	255	5
1,000	502	2
2,000	1,013	13
3,000	1,510	10
4,000	2,029	29
5,000	2,533	33
6,000	3,009	9
7,000	3,516	16
8,000	4,034	34
9,000	4,538	38
10,000	5,067	67

## Kerrich's results plotted



Instead of getting closer, the numbers of heads and tails are getting farther apart

## Repeating the experiment 50 times




This is not a fluke – instead, it occurs systematically and consistently in repeated simulated experiments

## Where's the law of averages?

- So where's the law of averages?
- Well, the law of averages does **not** say that as  $n$  increases the number of heads will be close to the number of tails
- What it says instead is that, as  $n$  increases, the average number of heads will get closer and closer to the long-run average (in this case, 0.5)
- The technical term for this is that the sample average, which is an estimate, *converges* to the *expected value*, which is a parameter

## Repeating the experiment 50 times, Part II





## Trends in Kerrich's experiment

- There are three very important trends going on in this experiment
- These trends can be observed visually from the computer simulations or proven via the binomial distribution
- We'll work with both approaches so that you can get a sense of how they both work and reinforce each other
- Before we do so, I'll introduce two additional, important facts about the binomial distribution: its mean (expected value) and standard deviation

## The expected value of the binomial distribution

- If we flip a coin once, how many heads do we expect (i.e., what is the expected value)? How about twice? Three times?  $n$  times?
- In each of those situations, what do we expect the average to be?

Tosses	1	2	3	...	$n$
Expected # of heads	0.5	1	1.5	...	$np$
Expected average	0.5	0.5	0.5	...	$p$

- The expected value of the binomial distribution is  $np$
- The expected value of a sample proportion  $\hat{p}$  (the average of a binomial distribution) is  $p$

## The standard error of the mean

- It can be shown that the standard deviation of the binomial distribution is

$$\sqrt{np(1-p)}$$

- What will the variability (standard deviation) be for one flip?  
Two?  $n$ ?
- What about the variability of the average?

Tosses	1	2	3	...	$n$
SD (# of heads)	0.5	0.71	0.87	...	$\sqrt{np(1-p)}$
SD (average)	0.5	0.35	0.29	...	$\sqrt{p(1-p)/n}$

## Standard errors

- Note that, as  $n$  goes up, the variability of the # of heads goes up, but the variability of the average goes down – just as we saw in our simulation
- Indeed, the variability goes to 0 as  $n$  gets larger and larger – this is the law of averages
- The standard deviation of the average is given a special name in statistics to distinguish it from the sample standard deviation of the data
- The standard deviation of the average is called the *standard error*
- The term *standard error* can also be applied to mean the variability of any estimate, to distinguish it from the variability of individual tosses or people

## The square root law

- The relationship between the variability of an individual (toss) and the variability of the average (of a large number of tosses) is a very important relationship, sometimes called the *square root law*:


$$SE = \frac{SD}{\sqrt{n}},$$

where SE is the standard error of the mean and SD is the standard deviation of an individual (toss)

- We saw that this is true for tosses of a coin, but it is in fact true for all averages
- Once again, we see this phenomenon visually in our simulation results

## The distribution of the mean

Finally, let's look at the distribution of the mean by creating histograms of the mean in our simulation



# The central limit theorem

- In summary, there are three very important phenomena going on here concerning the sampling distribution of the sample average:
  - #1 The expected value is always equal to the population average
  - #2 The standard error is always equal to the population standard deviation divided by the square root of  $n$
  - #3 As  $n$  gets larger, the sampling distribution looks more and more like the normal distribution
- Furthermore, these three properties of the sampling distribution of the sample average hold for **any distribution** – not just the binomial

## The central limit theorem (cont'd)

- This result is called the *central limit theorem*, and it is one of the most important, remarkable, and powerful results in all of statistics
- In the real world, we rarely know the distribution of our data
- But the central limit theorem says: we don't have to





## The central limit theorem (cont'd)

- Furthermore, as we have seen, knowing the mean and standard deviation of a distribution that is approximately normal allows us to calculate anything we wish to know with tremendous accuracy – and the sampling distribution of the mean is always approximately normal
- The only caveats:
  - Observations must be independently drawn from and representative of the population
  - The central limit theorem applies to the sampling distribution of the mean – not necessarily to the sampling distribution of other statistics
  - How large does  $n$  have to be before the distribution becomes close enough in shape to the normal distribution?

## How large does $n$ have to be?



- Rules of thumb are frequently recommended that  $n = 20$  or  $n = 30$  is “large enough” to be sure that the central limit theorem is working
- There is some truth to such rules, but in reality, whether  $n$  is large enough for the central limit theorem to provide an accurate approximation to the true sampling distribution depends on how close to normal the population distribution is
- If the original distribution is close to normal,  $n = 2$  might be enough
- If the underlying distribution is highly skewed or strange in some other way,  $n = 50$  might not be enough

## Example #1




## Example #2

Now imagine an urn containing the numbers 1, 2, and 9:




## Example #2 (cont'd)



## Example #3

- Weight tends to be skewed to the right (far more people are overweight than underweight)
- Let's perform an experiment in which the NHANES sample of adult men is the population
- I am going to randomly draw twenty-person samples from this population (*i.e.* I am re-sampling the original sample)

## Example #3 (cont'd)



## Why do so many things follow normal distributions?

- We can see now why the normal distribution comes up so often in the real world: any time a phenomenon has many contributing factors, and what we see is the average effect of all those factors, the quantity will follow a normal distribution
- For example, there is no one cause of height – thousands of genetic and environmental factors make small contributions to a person's adult height, and as a result, height is normally distributed
- On the other hand, things like eye color, cystic fibrosis, broken bones, and polio have a small number of (or a single) contributing factors, and do not follow a normal distribution



# Summary

- Central limit theorem:
  - The expected value of the average is always equal to the population average
  - $SE = SD/\sqrt{n}$
  - As  $n$  gets larger, the sampling distribution looks more and more like the normal distribution
- Generally speaking, the sampling distribution looks pretty normal by about  $n = 20$ , but this could happen faster or slower depending on the population and how skewed it is