

Likelihood Theory and Extensions (BIOS:7110)  
Breheny

Assignment 9

Due: Monday, November 11

1. *Score test with nuisance parameters.* This problem consists of proving the following theorem, which was stated in our lecture on score, Wald, and likelihood ratio approaches:

**Theorem:** If (A)-(C) hold and  $\boldsymbol{\theta}_0 = \boldsymbol{\theta}_1^*$ , then

$$\mathbf{u}_1(\hat{\boldsymbol{\theta}}_0)^\top \mathcal{I}_{11}^n(\hat{\boldsymbol{\theta}}_0) \mathbf{u}_1(\hat{\boldsymbol{\theta}}_0) \xrightarrow{d} \chi_r^2.$$

In this theorem, you may make use of the following lemma, which you do not need to prove (its proof is essentially identical to the case for the unrestricted MLE  $\hat{\boldsymbol{\theta}}$ ).

**Lemma:** If (A)-(C) hold and  $\boldsymbol{\theta}_0 = \boldsymbol{\theta}_1^*$ , then

$$\sqrt{n}(\hat{\boldsymbol{\theta}}_2(\boldsymbol{\theta}_0) - \boldsymbol{\theta}_2^*) = \mathcal{J}_{22}^{-1}(\boldsymbol{\theta}^*) \frac{1}{\sqrt{n}} \mathbf{u}_2(\boldsymbol{\theta}^*) + o_p(1),$$

where  $\mathbf{u} = (\mathbf{u}_1^\top \mathbf{u}_2^\top)^\top$  describes the partitioned score vector,  $\mathcal{J}_{22}$  is the lower right block of the Fisher information for a single observation, and here the final  $o_p(1)$  term is a  $(d-r) \times 1$  vector, all of whose elements are converging in probability to zero.

- (a) Take a Taylor series expansion of  $\frac{1}{\sqrt{n}} \mathbf{u}(\boldsymbol{\theta})$  about  $\boldsymbol{\theta}^*$ , evaluated at  $\hat{\boldsymbol{\theta}}_0$ .
  - (b) Show that, under the null,  $\frac{1}{\sqrt{n}} \mathbf{u}_1(\hat{\boldsymbol{\theta}}_0) = \mathbf{A} \frac{1}{\sqrt{n}} \mathbf{u}(\boldsymbol{\theta}^*) + o_p(1)$ , where  $\frac{1}{\sqrt{n}} \mathbf{u}_1(\hat{\boldsymbol{\theta}}_0)$  is the first component of the expansion in part (a) and  $\mathbf{A}$  is a matrix for you to determine.
  - (c) Using the result from (b), prove the above theorem.
2. *Projection matrices and the  $\chi^2$  distribution.* Suppose that  $\mathbf{z} \sim N(\mathbf{0}, \mathbf{I})$  and that  $\mathbf{P}$  is a symmetric projection matrix (i.e., that  $\mathbf{P} = \mathbf{P}\mathbf{P}$ ). Show that  $\mathbf{z}^\top \mathbf{P} \mathbf{z} \sim \chi_r^2$ , where  $r$  is the rank of  $\mathbf{P}$ . (you may wish to consult assignment 3 for some related results on projection matrices)
  3. *Score, Wald, and likelihood ratio tests for the rate parameter of the Gamma distribution.* This is a continuation of the problem, “Quadratic approximation for the Gamma distribution”. In that problem, you derived the score and information, and simulated a random sample from the Gamma distribution. Now, implement score, Wald, and likelihood ratio approaches for carrying out inference regarding the rate parameter  $\beta$ . We discussed these approaches conceptually in class; this problem ensures that you understand all the specifics well enough to program them. Specifically, turn in a separate .R file that defines the following six functions (where  $\mathbf{x}$  is the data, `level` is the confidence level, and `b0` is the hypothesized value of  $\beta$ ):

- `wald_ci(x, level=0.95)`
- `wald_test(x, b0=1)`
- `score_ci(x, level=0.95)`
- `score_test(x, b0=1)`

- `lr_ci(x, level=0.95)`
- `lr_test(x, b0=1)`

In other words, after sourcing the file you turn in, I should be able to run `score_ci(x)` and get a 95% confidence interval, but I should also be able to run `score_ci(x, 0.9)` and get a 90% confidence interval. The `_ci` functions should return a vector of length 2 (the lower and upper confidence limits), while the `_test` functions should return a vector of length 1 (the  $p$ -value).

Your `.R` file can define other functions (for example, you will likely want an `mle()` function, since many of these methods require the MLE), but it should not execute anything or load any packages.